

This document is reference information about the products.  
Please confirm a document attached to a product definitely.

# Medipure N Class-2 pharmaceutical product Anti-allergy Agent

Medipure N is a medicine you take orally to treat itching in sensitive areas. It is a peach mint flavored chewable tablet that can be taken immediately without water, even in situations where it is difficult to manage the problem, such as when you are out and about.

## ⚠ PRECAUTIONS

### ⊗ DON'Ts

**[Failure to follow these precautions may worsen the clinical condition or cause adverse reactions / incidents.]**

**1. Do not use if**

- (1) you have experienced allergic reactions to this medicine or its ingredients.
- (2) you have phenylketonuria (because this medicine contains an L-phenylalanine compound.)

**2. Do not take the following medicines while taking this medicine:**

Other medicines for allergies, internal medicines containing antihistamines, etc. (cold medicine, expectorant, internal medicine for rhinitis, medicine for motion sickness, etc.)

**3. Do not drive or operate machinery after taking this medicine** (as it may cause drowsiness)

**4. Avoid prolonged use**

**5. Avoid alcohol while taking this medicine**



### CONSULT

**1. Consult a doctor, pharmacist, or registered retail person before use if**

- (1) you are under the care of a doctor
- (2) you are pregnant or suspected of being pregnant
- (3) you are breastfeeding
- (4) you are elderly
- (5) you have ever had an allergic reaction to a medical product, etc.
- (6) you have following symptom: dysuria
- (7) you have been diagnosed with the following disease: glaucoma

**2. Discontinue use immediately and consult a doctor, pharmacist or registered retail person showing this package insert if any of the following symptoms occur after the medication, since it can be an adverse reaction.**

Affected area	Symptoms
Skin	rash/redness, itching
Gastrointestinal system	nausea and vomiting, loss of appetite, abdominal pain
Neuropsychiatric system	headache, light-headed feeling
Urinary system	dysuria

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid:

Condition	Symptoms
Shock (anaphylaxis)	Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after use.
Hepatic function failure	Symptoms, such as fever, itching, rash, jaundice (skin and sclera of the eyes turn yellow), brown urine, general malaise, loss of appetite, etc. may occur.
Thrombocytopenia	Symptoms, such as nosebleeds, bleeding from the gums and bruises, caused by reduced counts of platelets which is a component of the blood may occur.

3. The following symptoms may occasionally occur after taking the medication. If such symptoms persist or worsen, discontinue use and consult a doctor, pharmacist, or registered retail person showing this package insert.

Dry mouth, drowsiness

4. If the symptoms are not alleviated after 5 to 6 days of treatment, discontinue use of medication, and consult a doctor, pharmacist or registered retail person showing this package insert.

#### INDICATIONS

Itching due to eczema and/or rash, rhinitis, urticaria

#### DOSAGE AND ADMINISTRATION


For adults ( $\geq 15$  years of age), take 1 tablet at a time twice daily, in the morning and evening, by chewing or dissolving it in the mouth.

Do not give to children under 15 years of age.

#### <Precautions of Dosage and Administration>

- (1) Follow the dosage and administration limits strictly.
- (2) How to remove the tablets from blisters:

Press the blister with fingertips to break the aluminum foil on the reverse side and tip the tablet out before swallowing it, as shown in the right illustration. (Swallowing a tablet without removing the PTP sheet by mistake may cause unexpected problems such as the PTP sheet sticking to the mucous membrane of the throat.)



#### Ingredient and Purpose


The daily dose for adults (two tablets) contains:

Ingredient	Amount	Purpose
Mequitazine	4 mg	Reduces skin itching and rhinitis effectively through its anti-allergic effect

D-Mannitol, magnesium aluminometasilicate, hydroxypropylcellulose, tannic acid, sucralose, crospovidone, aspartame (L-phenylalanine), citric acid hydrate, l-menthol, magnesium stearate and aromatics are included as additives.

#### PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place, away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer to another container, in order to avoid misuse and maintain its quality.
- (4) Do not use after the expiration date.



Distributed by

**TANABE PHARMA CORPORATION**

3-2-10, Doshomachi, Chuo-ku, Osaka

Manufactured and distributed by

**Teika Pharmaceutical Co., Ltd.**

1-3-27, Arakawa, Toyama